

Zootaxa 2018 vol.4527 N1, pages 105-123

---

# Identification of *Ceriodaphnia dana*, 1853 (Crustacea: Cladocera) taxa from European Russia based on ehippial morphology

Kotov A., Ibragimova A., Neretina A.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

---

## Abstract

Copyright © 2018 Magnolia Press. Over the last decades significant progress was achieved in the investigations of water fleas (Crustacea: Cladocera). Their morphology, taxonomy and biogeography can be considered as well-studied, but still there are genera almost ignored by taxonomists. The genus *Ceriodaphnia* Dana, 1853 (Cladocera: Daphniidae) belongs to such problematic groups. From previous publications, it is obvious that different taxa from the genus have a very different morphology of their ehippia. Here, we study ehippium morphology in six common taxa from central-northern European Russia (*Ceriodaphnia megops* Sars, 1862; *C. laticaudata* P.E. Müller, 1867; *C. rotunda* (Straus, 1820) sensu Sars, 1862; *C. quadrangula* (O.F. Müller, 1785); *C. pulchella* Sars, 1862; *C. reticulata* (Jurine, 1820)) using light and scanning electron microscopy. A key to their identification based on ehippium morphology is proposed. This could be the starting point for revisions based on morphological characters with special emphasis on ehippia. Also, we propose that *Ceriodaphnia* taxa at least from the Holocene subfossil samples could be identified to the species group level according to structure of their ehippia. This approach to identification of *Ceriodaphnia* remains in sediments could improve palaeoecological reconstructions. However a global revision of the genus is still needed.

<http://dx.doi.org/10.11646/zootaxa.4527.1.9>

---

## Keywords

Cladocera, Ehippium, European Russia, Morphology, Taxonomy

## References

- [1] Abreu, M.J., Santos-Wisniewski, M.J., Rocha, O. & Orlando, TC. (2010) The use of PCR-RFLP to genetically distinguish the morphologically close species: *Ceriodaphnia dubia* Richard, 1894 and *Ceriodaphnia silvestrii* Daday, 1902 (Crustacea Cladocera). *Brazilian Journal of Biology*, 70, 121–124. <https://doi.org/10.1590/S1519-69842010000100016>
- [2] Alonso, M. (1996) *Fauna Iberica. Crustacea Branchiopoda*. Consejo Superior de Investigaciones Científicas, Madrid, 486 pp. Berner, D.B. (1985) Morphological differentiation among species in the *Ceriodaphnia cornuta* complex (Crustacea, Cladocera). *Verhandlungen der Internationalen Vereinigung für Theoretische und Angewandte Limnologie*, 22, 3099–3103. <https://doi.org/10.1080/03680770.1983.11897839>
- [3] Berner, D.B. (1986) Taxonomy of *Ceriodaphnia* (Crustacea: Cladocera) in U.S. Environmental Protection Agency Cultures. U.S. Environmental Protection Agency, Cincinnati, 34 pp.
- [4] Berner, D.B. (1987) Significance of head and carapace pores in *Ceriodaphnia* (Crustacea, Cladocera). *Hydrobiologia*, 145, 75–84. <https://doi.org/10.1007/BF02530267>

- [5] Berner, D.B. (1992) Restricted distribution of *Ceriodaphnia quadrangula* in North America. XXV SIL International Congress, Memorias, 1, 50.
- [6] Berner, D.B. & Rakhmatullaeva, G. (2001) A new species of *Ceriodaphnia* from Uzbekistan and Kazakhstan. *Hydrobiologia*, 442, 29–39. <https://doi.org/10.1023/A:1017586106850>
- [7] Bottrell, H.H. & Newsome, T. (1976) Body length, size and identification of the ephippia of *Ceriodaphnia megalops* and *Ceriodaphnia pulchella*. *Freshwater Biology*, 6, 405–408. <https://doi.org/10.1111/j.1365-2427.1976.tb01626.x>
- [8] Chatterjee, T., Kotov, A.A., Van Damme, K., Chandrasekhar, S.V.A. & Padhye, S. (2013) An annotated checklist of the Cladocera (Crustacea: Branchiopoda) from India. *Zootaxa*, 3667 (1), 1–89. <https://doi.org/10.11646/zootaxa.3667.1.1>
- [9] Deevey, E.S. (1969) Cladoceran populations of Rogers Lake during late- and postglacial time. *Verhandlungen der Internationalen Vereinigung für Theoretische und Angewandte Limnologie*, 17, 56–63.
- [10] Elías-Gutiérrez, M., Martínez Jerónimo, F., Ivanova, N.V., Valdez Moreno, M. & Hebert, P.D.N. (2008) DNA barcodes for Cladocera and Copepoda from Mexico and Guatemala, highlights and new discoveries. *Zootaxa*, 1839, 1–42.
- [11] Frey, D.G. (1959) The taxonomic and phylogenetic significance of the head pores of the Chydoridae (Cladocera). *Internationale Revue der gesamten Hydrobiologie und Hydrographie*, 44, 27–50. <https://doi.org/10.1002/iroh.19590440104>
- [12] Frey, D.G. (1960) The ecological significance of Cladocera remains in lake sediments. *Ecology*, 41, 684–699. <https://doi.org/10.2307/1931802>
- [13] Frey, D.G. (1962) Cladocera from the Eemian interglacial of Denmark. *Journal of Paleontology*, 36, 1133–1154. Available from: <https://www.jstor.org/stable/1301322> (Accessed 3 Dec. 2018)
- [14] Glagolev, S.M. (1983) A structure of ephippium surface in Daphniidae (Crustacea, Cladocera) by the data of scanning electron microscopy. *Zoologicheskii Zhurnal*, 62, 1422–1424.
- [15] Goulden, C.E. (1968) The systematics and evolution of the Moinidae. *Transactions of the American Philosophical Society Held at Philadelphia, New Series*, 58 (6), 1–101. <https://doi.org/10.2307/1006102>
- [16] Greenwood, T.L., Green, J.D. & Chapman, M.A. (1991) New Zealand *Ceriodaphnia* species: identification of *Ceriodaphnia dubia* Richard, 1894 and *Ceriodaphnia* cf. *pulchella* Sars, 1862. *New Zealand Journal of Marine and Freshwater Research*, 25, 283–288. <https://doi.org/10.1080/00288330.1991.9516480>
- [17] Hegna, T.A. & Kotov, A.A. (2016) Ephippia belonging to *Ceriodaphnia* Dana, 1853 (Cladocera: Anomopoda: Daphniidae) from the Lower Cretaceous of Australia. *Palaeontologia Electronica*, 19.3.40A, 1–9. <https://doi.org/10.26879/667>
- [18] Hofmann, W. (1984) Postglacial morphological variation in *Bosmina longispina* Leydig (Crustacea, Cladocera) from the Grosser Ploner See (North Germany) and its taxonomic applications. *Zeitschrift für Zoologische Systematik und Evolutionsforschung*, 22, 294–301. <https://doi.org/10.1111/j.1439-0469.1984.tb00664.x>
- [19] Hudec, I. (2010) Anomopoda, Ctenopoda, Haplopoda, Onychopoda (Crustacea: Branchiopoda). *Fauna Slovenska III. VEDA, Bratislava*, 496 pp.
- [20] Jaksch, H. (1992) Vergleichende rasterelektronenmikroskopische Untersuchungen an Daphniidae (Cladocera; Crustacea) Österreichs (Burgeländischer Seewinkel, Augebiete Östlich von Wien); Erfassung der Ephippial- und Revision der Carapaxskulpturen. Ph.D. Thesis, Universität Wien, Wien. Available from: [https://reflex.at/~biolab/Unterseiten/Diss/Titel\\_Inhalt.htm](https://reflex.at/~biolab/Unterseiten/Diss/Titel_Inhalt.htm) (accessed 6 November 2018)
- [21] Kořínek, V. (2002) Cladocera. In: Fernando, C.H. (Ed.), *A guide to tropical freshwater zooplankton*. Backhuys Publishers, Leiden, pp. 69–122.
- [22] Korosi, J.B. & Smol, J.P. (2012) An illustrated guide to the identification of cladoceran subfossils from lake sediments in northeastern North America: part 1—the Daphniidae, Leptodoridae, Bosminidae, Polyphemidae, Holopedidae, Sididae, and Macrothricidae. *Journal of Paleolimnology*, 48, 571–586. <https://doi.org/10.1007/s10933-012-9632-3>
- [23] Korovchinsky, N.M. (1995) *Ceriodaphnia* (Daphniidae). In: Alexeev V.R. (Ed.), *Key book for freshwater invertebrates of Russia and surrounding territories*. Vol. 2. Crustacea. KMK, Moscow, pp. 47–48.
- [24] Korovchinsky, N.M. (2004) Cladocerans of the order Ctenopoda of the world fauna (morphology, systematics, ecology, biogeography). KMK Press, Moscow, 410 pp. [in Russian]
- [25] Kotov, A.A. (2013) *Morphology and phylogeny of Anomopoda* (Crustacea: Cladocera). KMK, Moscow, 638 pp. [in Russian with English abstract]
- [26] Kotov, A.A. & Bekker, E.I. (2016) Cladocera: family Eurycercidae (Branchiopoda: Cladocera: Anomopoda). In: Dumont, H.J. (Ed.), *Identification guides to the plankton and benthos of inland waters*. Vol. 25. Backhuys Publishers, Leiden & Margraf Publishers, Weikersheim, pp. 89.
- [27] Kotov, A.A., Sinev, A.Y., Glagolev, S.M. & Smirnov, N.N. (2010) Water fleas (Cladocera). In: Alexeev, V.R. & Tsalolokhin, S.Y. (Eds.), *Key book for zooplankton and zoobenthos of fresh waters of European Russia*. KMK, Moscow, pp. 151–276.

- [28] Lampert, W. (2011) *Daphnia*: development of a model organism in ecology and evolution. *Excellence in Ecology*, 21, 1–250.
- [29] Lu, K.H. (2001) Morphology and ultrastructure of the ephippia of some Cladocera (Crustacea). *Acta zoologica Sinica*, 47, 565–572.
- [30] Mergeay, J., Verschuren, D. & De Meester, L. (2005) *Daphnia* species diversity in Kenya, and a key to the identification of their ephippia. *Hydrobiologia*, 542, 261–274. <https://doi.org/10.1007/s10750-004-4952-6>
- [31] Moritz, C. (1988) Die Verteilung der Ephippien von *Bosmina longirostris* und *Ceriodaphnia pulchella* im Sediment des Piburger Sees (Ötztal, Tirol) (Cladocera, Crustacea). *Berichte des naturwissenschaftlichen-medizinischen Verein Innsbruck*, 75, 91–107.
- [32] Popova, E.V., Petrusek, A., Kořínek, V., Mergeay, J., Bekker, E.I., Karabanov, D.P., Galimov, Y.R., Neretina, T.V., Taylor, D.J. & Kotov, A.A. (2016) Revision of the Old World *Daphnia* (Ctenodaphnia) similis group (Cladocera: Daphniidae). *Zootaxa*, 4161 (1), 1–40. <https://doi.org/10.11646/zootaxa.4161.1.1>
- [33] Sharma, P. (2014) Morphological and molecular identification of three *Ceriodaphnia* species (Cladocera: Daphniidae) from Australia. *Advances in Zoology*, 2014, 25813. <https://doi.org/10.1155/2014/258134>
- [34] Sharma, P. & Kotov, A.A. (2013) Molecular approach to identify sibling species of the *Ceriodaphnia cornuta* complex (Cladocera: Daphniidae) from Australia with notes on the continental endemism of this group. *Zootaxa*, 3702 (1), 79–89. <https://doi.org/10.11646/zootaxa.3702.1.5>
- [35] Smirnov, N.N. (1992) The Macrothricidae of the world. Guides to the identification of the microinvertebrates of the Continental Waters of the world. SPB Academic Publishing, The Hague, 143 pp.
- [36] Smirnov, N.N. (1996) Cladocera: the Chydorinae and Sayciinae (Chydoridae) of the world. Guides to the identification of the microinvertebrates of the Continental Waters of the world. SPB Academic Publishing, Amsterdam, 197 pp.
- [37] Smirnov, N.N. (2010) Historic ecology of lake zoocenoses. KMK, Moscow, 225 pp. [in Russian]
- [38] Smirnov, N.N. (2017) Physiology of the Cladocera. 2nd Edition. Academic Press, London, 418 pp.
- [39] Szeroczyńska, K. & Sarmaja-Korjonen, K. (2007) Atlas of subfossil Cladocera from Central and Northern Europe. Friends of the Lower Vistula Society, Świecie, 83 pp.
- [40] Van Damme, K. & Kotov, A.A. (2016) The fossil record of the Cladocera (Crustacea: Branchiopoda): Evidence and hypotheses. *Earth-Science Reviews*, 163, 162–189. <https://doi.org/10.1016/j.earscirev.2016.10.009>
- [41] Vandekerkhove, J., Declerck, S., Vanhove, M., Brendonck, L., Jeppesen, E., Conde Porcuna, J.M. & De Meester, L. (2004) Use of ephippial morphology to assess richness of anomopods: potentials and pitfalls. *Journal of Limnology*, 63 (Supplement 1), 75–84. <https://doi.org/10.4081/jlimnol.2004.s1.75>
- [42] Zawiska, I., Zawisza, E., Wojewódka, M. & Sinev, A.Y. (2016) Exploring the world of micro sculptures-subfossil Cladocera remains under the SEM. *Advances in Oceanography and Limnology*, 7 (2), 177–183. <https://doi.org/10.4081/aiol.2016.6218>